

ON THE NATURE OF THE FAST MOVING STAR S2 IN THE GALACTIC CENTER *

F. MARTINS ¹, S. GILLESSEN ¹, F. EISENHAEUER ¹, R. GENZEL ^{1,2}, T. OTT ¹, S. TRIPPE ¹

ABSTRACT

We analyze the properties of the star S2 orbiting the supermassive black hole at the center of the Galaxy. A high quality SINFONI H and K band spectrum obtained from coadding 23.5 hours of observation between 2004 and 2007 reveals that S2 is an early B dwarf (B0–2.5V). Using model atmospheres, we constrain its stellar and wind properties. We show that S2 is a genuine massive star, and not the core of a stripped giant star as sometimes speculated to resolve the problem of star formation so close to the supermassive black hole. We give an upper limit on its mass loss rate, and show that it is He enriched, possibly because of the presence of a magnetic field.

Subject headings: Stars: early-type — stars: fundamental parameters — Galaxy: center

1. INTRODUCTION

The central parsec of our Galaxy hosts a large population of young massive stars (Krabbe et al. 1995; Paumard et al. 2006). Their presence is puzzling since according to standard theories, star formation should not happen so close to the supermassive black hole SgrA*: the tidal forces are so large that any molecular cloud should be disrupted before being able to collapse (Morris 1993). For most of the young stars, the solution seems to be star formation in dense accretion disks as witnessed by the two counter-rotating flat stellar structures around SgrA* (Levin & Beloborodov 2003; Genzel et al. 2003; Paumard et al. 2006). However, this attractive solution does not explain the so-called “S stars”, the group of objects located within one arcsecond (=0.038 pc) of the black hole. Their orbits are randomly oriented (Eisenhauer et al. 2005) and thus do not fit the accretion disk scenario. Spectroscopically identified as B stars (Ghez et al. 2003; Eisenhauer et al. 2005), their true nature as massive stars has been challenged: they have been proposed to be the hot, luminous cores of evolved red giant and/or AGB stars the envelope of which has been stripped by tidal interactions with SgrA* (Alexander 2005; Davies & King 2005). In that case, the problem of their formation process vanishes since they might very well have formed away from the hostile environment of the black hole before having migrated towards the Galactic Center. Such a process is too slow to explain the presence of short lived massive stars close to SgrA*. Hence, these S stars constitute a “paradox of youth” (Ghez et al. 2003).

In this letter, we analyze the properties of the brightest S-star – S2 – and show that it is a genuine massive star.

2. OBSERVATIONAL DATA AND MODELS

To constrain the physical parameters of S2, a high quality spectrum is needed. Since the observation of the S stars is time consuming, the S/N ratio obtained

in one night is usually limited to 10 at maximum. To increase this ratio, we have co-added all the S2 spectra obtained with SINFONI (Eisenhauer et al. 2003) in adaptive optics mode (0.0125''×0.025'') on the VLT since 2004. This corresponds to a total integration time of 23.5 hours in K band. Each spectrum was carefully extracted by selection of source and background pixels from which nebular contamination was removed. Note however that this is not such a critical problem for S2 which has a large radial velocity so that the position of the stellar absorption core is blueshifted beyond the nebular emission. We have determined the combined spectrum by a cross-correlation technique. First, the position of the Br γ line was estimated by a simple Gaussian fit to the line. From the such obtained radial velocities a first combined spectrum was calculated. This spectrum was then used to cross-correlate all individual spectra against it. This yielded better estimates for the radial velocities, which in turn were used to get a more accurate combined spectrum. This procedure was iterated until the result did not change anymore. The resulting spectrum shown in Fig. 1 has a S/N ratio of a few tens in the K band and a resolution of ~ 4000 . The H band spectrum is noisier due to shorter observation time and higher extinction.

The main K band lines (He I 2.112 μm and Br γ) are clearly identified. But we also detect several weak He I lines at 2.149, 2.161 and 2.184 μm . Similarly, one He I and three H I lines are identified for the first time in the H band ³. We do not see any trace of He II 2.189 μm . With the current S/N ratio, this line would be detected if it had a depth of about 1% of the continuum level.

For the quantitative analysis of this spectrum, we used the atmosphere code CMFGEN (Hillier & Miller 1998). It allows the computation of non-LTE atmosphere models including winds and line blanketing and is appropriate for hot massive stars. A description of the code is given in Hillier & Miller (1998), and we refer to Martins et al. (2007) for a summary of the procedure used to build the models. To derive the parameters of interest, we have run models with: $19000 < T_{\text{eff}} < 30000$ K, $3.0 < \log g < 4.75$, $0.1 < \text{He}/\text{H} < 2.0$ and $10^{-8} < \dot{M} < 10^{-5.5} \text{ M}_{\odot} \text{ yr}^{-1}$. These models and synthetic spectra included, in addition to H and He, C, N, O, Si, S and Fe. We adopted

³ These lines were seen by Eisenhauer et al. (2005) in the average spectrum of several stars, but not in individual spectra.

*BASED ON OBSERVATIONS COLLECTED AT THE ESO VERY LARGE TELESCOPE (PROGRAMS 075.B-0547, 076.B-0259, 077.B-0503, 078.B-0520 AND 179.B-0261)

Electronic address: martins@mpe.mpg.de

¹ Max Planck Institute of Extraterrestrial Physics, Postfach 1312, D-85741, Garching, Germany

² Department of Physics, University of California, CA 94720, Berkeley, USA




FIG. 1.— Spectrum of S2 after addition of all the SINFONI observations since 2004. The main lines are indicated. the solar abundances of Grevesse & Sauval (1998).

3. THE NATURE OF S2: A GENUINE B STAR

3.1. Spectroscopic classification

The spectrum shown in Fig. 1 is typical of an early B star (see also Ghez et al. 2003). Inspection of the atlases of Wallace & Hinkle (1997) and Hanson et al. (2005) shows that for late O stars, He II 2.189 μm is expected. We do not detect this line in spite of the good enough S/N ratio. For stars with spectral types later than B3, the He I lines vanish. Since we detect several of them, we can safely argue that S2 has a spectral type between B0 and B2.5. Fig. 12 of Hanson et al. (2005) shows that in supergiants and giants, Br γ is clearly separated from He I 2.161 μm , while for dwarfs Br γ is broad enough to merge with the He I line. This latter morphology is similar to what we observe for S2 (see below for a quantitative comparison). Hence, in addition to being spectroscopically identified as an early B star, we can conclude from these qualitative arguments that *S2 is also a dwarf and not a supergiant/giant.*

3.2. A mass estimate

The key question we want to answer here is the exact nature of S2. Spectroscopically, S2 is unambiguously an early B dwarf. However, this does not necessarily mean that S2 is a young, massive star. It could be the core of an older, evolved star which would have lost its envelope through tidal interaction with SgrA* but would still spectroscopically look like a B star. In that case, the star could have formed far away from SgrA* before being dragged to its proximity and experiencing an envelope stripping. There would be no paradox of youth.

Recently, Davies & King (2005) and Dray et al. (2006) modeled this stripping process as well as the subsequent evolution of the remaining core in the vicinity of SgrA*. They showed that, under certain condition on the IMF and the capture rate, the population of S stars could be explained by tidal stripping. The argument is however based on the number of stars predicted at the position of the S stars in the HR diagram. Expressed differently, the conclusion comes from the fact that the modeled stripped

TABLE 1
PHYSICAL PROPERTIES DERIVED FROM OUR QUANTITATIVE ANALYSIS FOR DIFFERENT T_{eff} APPROPRIATE FOR B0–2.5V STARS.

T_{eff} [K]	19000	22000	25000	27000	30000
$\log \frac{L}{L_{\odot}}$	4.30	4.45	4.60	4.65	4.80
$\log g$	3.80	4.00	4.00	4.00	4.25
$\log g_{\text{min}}$	3.55	3.72	3.77	3.89	3.86
R [$R_{\odot}$]	13.1	11.6	10.7	9.7	9.4
R_{min} [$R_{\odot}$]	11.4	10.1	9.3	8.5	7.3
M_{min} [$M_{\odot}$]	16.8	19.5	18.6	20.5	14.1
He/H [#]	1.20	0.50	0.45	0.55	0.80
He/H $_{\text{min}}$ [#]	0.85	0.25	0.25	0.30	0.30
$\dot{M}$ [$10^{-7} M_{\odot} \text{ yr}^{-1}$]	<3	<3	<3	<3	<3

stars can reach the luminosities and effective temperatures of B stars. But strictly speaking, this does not exclude that the S stars are genuine massive stars, which would lie at the same position in the HR diagram.

For that, the only parameter which can unambiguously be used is the stellar mass. The most massive stars able to experience envelope stripping are AGB stars. According to stellar evolution, such objects are the evolved descendants of main sequence stars with $M < 8 M_{\odot}$. The core of such objects is actually much less massive: Table 4 of Forestini & Charbonnel (1997) shows that their mass is lower than $1 M_{\odot}$. In the scenario explored by Dray et al. (2006), only stars having lost at least 99% of their envelope could explain the S stars. Thus, their mass should be at most $\sim 1 M_{\odot}$.

To estimate the mass of S2, we can rely on gravity and radius: $M = gR^2/G$ where g is the gravity, R the radius and G the gravitational constant. Gravity can be derived from the shape of Br γ , while radius is straightforwardly obtained from the knowledge of effective temperature and luminosity. Below, we explain how we proceeded to constrain these parameters.

Effective temperature. T_{eff} is usually derived from the ratio of lines from two consecutive ionization states: He I / He II lines for O stars and Si III / Si IV for early B stars. Unfortunately, S2 does not show He II and Si lines in the K band. Hence, we can only rely on semi-quantitative arguments to estimate T_{eff} . We have shown above that S2 is a B0V to B2.5V star. According to the recent studies of Dufton et al. (2006) and Trundle et al. (2007), such stars have $19000 < T_{\text{eff}} < 30000$ K. We have thus run models for $T_{\text{eff}} = 19, 22, 25, 27$ and 30 kK.

Luminosity. To estimate the star's luminosity, for each T_{eff} we adjusted $\log \frac{L}{L_{\odot}}$ in our models to match the absolute K band magnitude of S2. This magnitude is defined as $M_K = m_K - A_K - DM$ with $m_K = 14.0$ the observed magnitude of S2 (Paumard et al. 2006), $A_K = 2.25$ the K band extinction at the position of S2 (Schödel et al. 2007) and $DM = 14.50$ the distance modulus for a distance to the Galactic Center of 7.94 kpc (Eisenhauer et al. 2003). The resulting absolute K magnitude of S2 is -2.75. The range of values we obtain for $\log \frac{L}{L_{\odot}}$ is 4.30–4.80. In practice, the uncertainty of ± 0.2 on A_K and 0.5 kpc on the distance to the Galactic Center translate into an uncertainty of about 0.12 in $\log \frac{L}{L_{\odot}}$.

Radius. The radius of the star is simply derived from $L = 4\pi\sigma_B R^2 T_{\text{eff}}^4$ (σ_B being the Boltzmann constant).




FIG. 2.— Determination of $\log g$ for the case $T_{\text{eff}} = 22000$ K. The black solid line is the observed $\text{Br}\gamma$ line. The dashed (dotted, dot-dashed) line is a model with $\log g = 4.0$ (3.00, 4.50). All other parameters are kept constant (especially $\text{He}/\text{H}=0.5$). The wavelength range used for the $\log g$ determination is indicated.

The minimum radius allowed by our study (corresponding also to the minimum mass of the star – see below) is estimated for a luminosity reduced by the uncertainty (0.12 dex). The values of R_{min} are given in Table 1.

Gravity. $\log g$ was constrained from the shape of $\text{Br}\gamma$. Repolust et al. (2005) showed that its wings (resp. absorption core) get broader (resp. weaker) when gravity increases. For large $\log g$, the blue wing merges with the He I 2.161 μm line as discussed previously. Fig. 2 shows how the line profile changes when $\log g$ increases from 3.0 to 4.5 for $T_{\text{eff}} = 22000$ K. The best models correspond to $\log g \sim 4.0$. A quantification of the goodness of the fit in the $\text{Br}\gamma$ region of the spectrum was performed by means of a χ^2 analysis. To minimize contamination by HeI, we used only the range 2.164–2.179 μm . Fig. 3 shows the result of this analysis and confirms that $\log g = 4.0$ is preferred. From this curve, we also derive a 3σ lower limit to $\log g$ ($\log g_{\text{min}}$). Table 1 shows the results for other T_{eff} . The values of $\log g$ we obtain are all typical of a dwarf star ($\log g \sim 4.0$, Trundle et al. 2007), confirming our previous “spectroscopic” findings.

Rotational velocity. We derived a projected rotational velocity of 100 ± 30 km s^{-1} . It best accounts for the shape of the He I 2.112 μm doublet: for larger $V \sin i$ the lines merge into a single component, for lower values, they are too separated. This value is commonly found for early B stars: according to Abt et al. (2002), $V \sin i = 127 \pm 8$ (108 ± 8) km s^{-1} for B0–2V (B3–5V) stars.

Table 1 summarizes, for each T_{eff} , the values of $\log g$ and radius. Since we want to test if S2 is the core of an AGB star, we also give the minimum mass of S2. All our estimates are larger than $14.1 M_{\odot}$. This is larger than the initial mass of AGB stars, and consequently is well above the mass of the core of such a star. From that, one can thus safely conclude that *S2 is not a stripped AGB star, and is really a genuine B star.*

4. MASS LOSS RATE AND HELIUM CONTENT

4.1. Mass loss rate




FIG. 3.— χ^2 as a function of $\log g$ for the $T_{\text{eff}} = 22000$ K model. The best fit model has $\log g = 4.0$, as also seen from Fig. 2.

Below a certain threshold, $\text{Br}\gamma$ is insensitive to $\dot{M}$. When $\dot{M}$ increases above this limit, the $\text{Br}\gamma$ absorption profile is progressively filled by emission. By determining this threshold, we can set an upper limit on the mass loss rate of S2 (see Fig. 4). In practice, the quantity we constrain is the wind density $\rho = \dot{M}/(4\pi R^2 v_{\infty})$ (where R is the stellar radius and v_{∞} the terminal velocity). Deriving a mass loss rates implies we know v_{∞} . Since the lines we observe in the S2 K band spectrum are insensitive to v_{∞} , we have to assume values. Terminal velocities of B dwarfs are difficult to estimate since the winds are usually weak and the spectra do not show P-Cygni lines or emission lines from which it is usually derived. Hence, we have adopted $v_{\infty} = 1000$ km s^{-1} which is appropriate for late O dwarfs (e.g. Bouret et al. 2003). If v_{∞} would be lower than 1000 km s^{-1} , estimated mass loss rates would be lower. Our assumptions thus lead to conservative upper limits of $\dot{M} \lesssim 3 \times 10^{-7} M_{\odot} \text{ yr}^{-1}$. This is lower than $\dot{M}$ required by Loeb (2004) to feed SgrA* without transport of angular momentum.

4.2. Helium content

The near infrared spectrum of S2 is dominated by H and He lines. We thus could constrain the He/H abundance ratio. Qualitatively, when this ratio increases, all He I lines have stronger absorption profiles, while H I lines get weaker. In practice, we used the He I 2.149 μm and He I 2.184 μm lines to constrain He/H since these lines are almost insensitive to microturbulence, which is not the case of He I 2.112 μm and He I 2.161 μm . Fig. 5 shows the changes in the He I line profiles when He/H is varied between 0.1 and 0.5 (model with $T_{\text{eff}} = 22000$ K). We find that the He/H ratio varies from about 0.45 at $T_{\text{eff}} = 25000$ K up to as high as 1.20 at $T_{\text{eff}} = 19000$ K. We proceeded as for $\log g$ (χ^2 analysis) to derive a 3σ lower limit on the value of He/H . All these lower limits are larger than 0.25, confirming that S2 is He enriched.

With $\text{He}/\text{H} > 0.25$, S2 falls into the category of the so-called “He rich” stars, a class of B stars with He/H between 0.3 and 10 (Smith 1996). Interestingly, these peculiar stars have a very narrow distribution of spectral




FIG. 4.— Effect of mass loss on Br γ . When $\dot{M}$ increases, Br γ starts to depart from a pure absorption to an emission profile. Characterizing the value of $\dot{M}$ for which this occurs provides an upper limit on the mass loss rate of S2.

types centered around B2, similar to what we find for S2 (B0–2.5V). The best studied He-rich star is σ OriE. The origin of its surface abundance pattern is explained by a combination of specific wind properties and magnetic field. The star should have a wind weak enough for ion decoupling to occur, i.e. the radiative acceleration, essentially due to metals, is not redistributed among the passive plasma (mainly H and He) because of a too low density and reduced Coulomb forces (Krtićka & Kubát 2001). In that case, Helium might accumulate at the surface of the star. But this extra Helium can remain on the surface only if turbulence, and its associated mixing effect, is suppressed. This calls for a magnetic field strong enough to freeze the stellar surface (Hunger & Groote 1999). Our upper limit on the mass loss rate of S2 is consistent with a relatively weak wind. In this scenario, chemical inhomogeneities (spots) are expected on the surface as a result of the interplay between the magnetic field geometry and the wind. We cannot test this prediction with our observations since we do not spatially resolve the stellar surface of S2. Similarly, no CNO abnormalities are expected in this model, which we cannot

test either due to the absence of CNO lines (observed and expected) in the near-IR spectrum of S2.

If the wind is strong enough to develop into a normal homogenous outflow, one might still observe a significant He enrichment if the star has a magnetic field and rotates fast enough. Indeed Maeder & Meynet (2005) predict that a strong (10^4 G) magnetic field can lead to solid body rotation of the star. This favors the diffusion of species by meridional circulation. As a consequence, a star can experience strong surface He enrichment: Fig. 10 of Maeder & Meynet (2005) shows that a $15 M_{\odot}$ star can reach a He mass fraction of 0.3 (around 0.12 in He/H) in 12 Myr. Contrary to the previous scenario, large N and reduced C abundances are also expected.

In view of the available models and of the currently known stellar properties, we thus tentatively propose that S2 is a magnetic star.




FIG. 5.— Determination of He/H. The black solid line is the observed spectrum of S2. The red dashed (blue dotted) line is a model with He/H=0.10 (0.50). $T_{\text{eff}} = 22000$ K and $\log g = 4.0$. The He rich model is required to fit the S2 spectrum.

We thank John Hillier for his help with CMFGEN, as well as Corinne Charbonnel and Ana Palacios for interesting discussions on AGB stars. We thank Andrea Ghez and an anonymous referee for useful comments.

REFERENCES

- Abt, H., et al., 2002, *ApJ*, 573, 389
 Alexander, T., 2005, *PhR*, 419, 65
 Bouret, J. C., et al., 2003, *ApJ*, 595, 1182
 Davies, M.B. & King, A.R., 2005, *ApJ*, 624L, 25
 Dray, L.M., et al., 2006, *MNRAS*, 372, 31
 Dufton, P.L., et al., 2006, *A&A*, 457, 265
 Eisenhauer, F., et al., 2003, *ApJ*, 597L, 121
 Eisenhauer, F., et al., 2003, *SPIE*, 4841, 1548
 Eisenhauer, F., et al., 2005, *ApJ*, 628, 246
 Figer, D.F., et al., 1999, *ApJ*, 506, 384
 Forestini, M. & Charbonnel, C., 1997, *A&AS*, 123, 241
 Genzel, R., et al., 2003, *ApJ*, 594, 812
 Ghez, A.M., et al., 2003, *ApJ*, 586L, 127
 Grevesse, N. & Sauval, A.J., 1998, *SSRv*, 85, 161
 Hanson, M.M., et al., 2005, *ApJS*, 161, 154
 Hillier, D.J., Miller, 1998, *ApJ*, 496, 407
 Hunger, K. & Groote, D., 1999, *A&A*, 351, 554
 Krabbe, A., et al., 1995, *ApJ*, 447L, 95
 Krtićka, J. & Kubát, J., 2001, *A&A*, 369, 222
 Levin, Y. & Beloborodov, A.M., 2003, *ApJ*, 590L, 33
 Loeb, A., 2004, *MNRAS*, 350, 725
 Martins, F. et al., 2007, *A&A*, 468, 233
 Meynet, G., Maeder, A., 2005, *A&A*, 440, 1041
 Morris, M., 1993, *ApJ*, 408, 496
 Paumard, T., et al., 2006, *ApJ*, 643, 1011
 Repolust, T. et al., 2004, *A&A*, 415, 349
 Repolust, T. et al., 2005, *A&A*, 440, 261
 Schödel, R., et al., 2007, *A&A*, 469, 125
 Smith, K.C., 1996, *Ap&SS*, 237, 77
 Trundle, C., et al., 2007, *A&A*, 471, 625
 Wallace, L., Hinkle, K., 1997, *ApJS*, 111, 445